

# Preliminary results from characterization of the Iberian Peninsula sturgeon based on analysis of the mtDNA cytochrome *b*

A. Almodóvar<sup>1</sup>, A. Machordom<sup>2</sup> and J. Suárez<sup>1</sup>

<sup>1</sup> IMIA, Madrid Regional Government, El Encín, PO Box 127, E-28800 Alcalá de Henares, Madrid, Spain. E-mail: ana.almodovar@imia.comadrid.es

<sup>2</sup> Museo Nacional de Ciencias Naturales, CSIC, José Gutiérrez Abascal 2, E-28006 Madrid, Spain

Received September 1999. Accepted June 2000.

## ABSTRACT

Historically, the Atlantic sturgeon *Acipenser sturio* L., 1758 was considered the only sturgeon species that inhabited rivers of the Iberian Peninsula. Nevertheless, in a recent paper, Garrido-Ramos *et al.* (1997) identified specimen EBD 8174 (museum collection of the Doñana Biological Station, Spain), from the Guadalquivir River, as the Adriatic sturgeon *Acipenser naccarii* Bonaparte, 1836, suggesting that both *A. sturio* and *A. naccarii* could be native to Spain. To test this theory, we compared partial mitochondrial cytochrome *b* (cyt *b*) gene sequences obtained from museum specimens of *A. sturio* that originated from the Iberian Peninsula, the Adriatic and the North Sea, as well as from live individuals of the Gironde River population in France. Specimens of *A. naccarii* and of the Siberian sturgeon *Acipenser baerii* Brandt, 1869 from fish farms were also included in the study. DNA from museum specimens was successfully amplified using the protocol of France and Kocher (1996) for DNA extraction from formalin-fixed and ethanol-preserved samples. Phylogenetic analysis was performed on partial cyt *b* gene sequences (over 402-bp and 245-bp). We identified specimen EBD 8174 as *A. sturio*, confirming the opinion that *A. sturio* is the only sturgeon species of the Iberian Peninsula. Further molecular analyses of museum specimens are needed for a description of the historical intraspecies genetic variation within *A. sturio*. This information is crucial for any future recovery plan for this species. Our comparison also showed interspecies sequence divergence ranging from 6.91 % (*A. sturio/A. naccarii*) to 7.43 % (*A. sturio/A. baerii*).

**Key words:** *Acipenser sturio*, *Acipenser naccarii*, *Acipenser baerii*, genetic variation, molecular analysis.

## RESUMEN

**Resultados preliminares de la caracterización del esturión de la península Ibérica mediante el análisis del gen citocromo *b***

Históricamente, el esturión atlántico *Acipenser sturio* L., 1758 ha sido considerado la única especie de esturión que vive en los ríos de la península Ibérica, siendo así demostrado por múltiples estudios morfológicos recientes. Sin embargo, un trabajo reciente identificó el ejemplar del río Guadalquivir EBD 8174 como *Acipenser naccarii* Bonaparte, 1836, sugiriendo que ambas especies de esturión podrían ser nativas de España. Para comprobar esta nueva teoría, se han analizado las secuencias del gen citocromo *b* mitocondrial en muestras de esturión atlántico preservadas en varios museos procedentes de la península Ibérica, el mar Adriático y el mar del Norte, así como material fresco del río Garona. Además, se han estudiado con fines comparativos varias muestras de *A. naccarii* y *Acipenser baerii* Brandt, 1869 procedentes de piscifactorías. Las muestras de museos fueron amplificadas con éxito usando un protocolo específico utilizado para material fijado en formol y preservado en alcohol. La muestra de esturión de España (EBD 8174) fue identificada como *A. sturio*. Este resulta-

do confirma que *A. sturio* es la única especie de esturión en la península Ibérica. De cualquier manera, antes de abordar los planes de recuperación, sería necesario realizar análisis moleculares complementarios para describir las posibles variaciones genéticas existentes entre poblaciones. La comparación de las secuencias de citocromo b de *A. sturio* con *A. naccarii* y *A. baerii* mostró valores de divergencia de 6,91 % y 7,43 % respectivamente.

**Palabras clave:** *Acipenser sturio*, *Acipenser naccarii*, *Acipenser baerii*, variación genética, análisis molecular.

## INTRODUCTION

The Atlantic sturgeon *Acipenser sturio* L., 1758, is native to Western Europe and historically was present in many large Spanish rivers (Holčík *et al.*, 1989). Moreover, *A. sturio* was considered the Iberian Peninsula's only sturgeon (Almaça, 1988; Doadrio, Elvira and Bernat, 1991; Elvira, Almodóvar and Lobón-Cerviá, 1991a, b; Elvira and Almodóvar, 1993; Pereira, 1995). The *A. sturio* population in the Guadalquivir River was intensively exploited for caviar beginning in the 1930s, and its population dropped dramatically from the 1960s (Elvira, Almodóvar and Lobón-Cerviá, 1991a,b).

Contrary to the prevailing scientific opinion, Garrido-Ramos *et al.* (1997) recently suggested that two sturgeon species could be native to Spain: not only *A. sturio*, but also the Adriatic sturgeon *Acipenser naccarii* Bonaparte, 1836, which was formerly recognized as endemic to the Adriatic Sea basin only (Tortonese, 1989; Birstein and Bemis, 1997). In addition to a revaluation of morphological data, Garrido-Ramos *et al.* (1997) claimed that their conclusion was supported by a molecular identification of two specimens, EBD 8173 and EBD 8174, from the collection of the Doñana Biological Station (Spain), as *A. naccarii*. However, three laboratories in different countries could not confirm these results because the paper of Garrido-Ramos *et al.* (1997) did not contain a description of the molecular methods used, and, therefore, its results could not be reproduced (Doukakis *et al.*, 2000). Doukakis *et al.* (2000) suggested that Garrido-Ramos *et al.* (1997) might have worked not with the authentic extracted DNA, but with a contaminant. Furthermore, new morphological studies of these two specimens, and a comparison of them with many museum sturgeon specimens from rivers of the Iberian Peninsula, are in disagreement with the conclusions of Garrido-Ramos *et al.* (1997) (Elvira and Almodóvar, 1999, 2000; Rincón, 2000).

In the meantime, data concerning the molecular taxonomy of *A. sturio* are scarce (Wirgin, Stabile and

Waldman, 1997; Birstein, Betts and DeSalle, 1998; Birstein and DeSalle, 1998; Ludwig and Kirschbaum, 1998). A molecular study of all available *A. sturio* specimens, including those kept in different museums, is crucial to any recovery programme for this species (Birstein and Doukakis, 2000; Ludwig *et al.*, 2000). Consequently, the goals of the present study were: (1) To elaborate a reliable method of DNA extraction from the museum (archival) specimens; (2) To identify and characterize by molecular methods the *A. sturio* specimens from the Iberian Peninsula and other European areas; (3) To determine the relationships between *A. sturio* and related sturgeon species.

## MATERIALS AND METHODS

### Sample collection

For DNA extraction, we used muscle tissue (from fresh, stuffed or ethanol-preserved specimens) from 7 specimens of *A. sturio*, 3 *A. baerii* and 3 *A. naccarii*, from different localities (table I). Fresh material was stored at -80 °C until DNA extraction.

### DNA extraction

Homogenization after freezing of fresh tissue samples in liquid nitrogen was followed by a standard procedure of phenol-dichloromethane extraction and alcohol precipitation (Wirgin *et al.*, 1997). Tissues from stuffed samples were incubated in 1 ml of shaking TE9 buffer (500 mM Tris, 20 mM EDTA, 10 mM NaCl, pH 9.0; Shiozawa *et al.*, 1992) at room temperature for 24 hours, with one change of the buffer. The TE9 buffer was discarded and the tissue samples were then minced and incubated for 5 hours at 55 °C in TE9 buffer containing 50 µl of 20 % sodium dodecyl sulfate (SDS) and 12.5 µl of 30 mg/ml Proteinase K and incubated at 55 °C during 5 hours. Thereafter, another 12.5 µl of 20 mg/ml Proteinase K was added and the tissue

Table I. List of sturgeon samples studied and cytochrome *b* regions sequenced

Sample	Species	Collection/Museum	Location of capture or farm	Preservation	Fragment of cyt <i>b</i>
S1	<i>A. sturio</i>	Doñana Biological Station, Seville EBD 8174, May 1975	Guadalquivir River, Alcalá del Río, Seville, Spain	Stuffed	155-bp
S2	<i>A. sturio</i>	Cemagref, Bourdeaux	Gironde River, France	Fresh	402-bp
S3	<i>A. sturio</i>	Cemagref, Bourdeaux	Gironde River, France	Fresh	402-bp
S4	<i>A. sturio</i>	Cemagref, Bourdeaux	Gironde River, France	Fresh	402-bp
S5	<i>A. sturio</i>	Cemagref, Bourdeaux	Gironde River, France	Fresh	402-bp
S6	<i>A. sturio</i>	Senckenberg Museum, Frankfurt SMF 2448	Adriatic Sea, Trieste, Italy	Ethanol	155-bp
S7	<i>A. sturio</i>	Senckenberg Museum, Frankfurt SMF 7637	Eider River, Nübbel, Germany	Ethanol	155-bp
B1	<i>A. baerii</i>	Doñana Biological Station, Seville EBD 8157, September 1995	Guadalquivir River, Coria del Río, Seville, Spain (released)	Fresh	402-bp
B2	<i>A. baerii</i>	Department of Animal Biology I Complutense University, Madrid, 1996	Duración River, Spain (released)	Fresh	402-bp
B3	<i>A. baerii</i>	Department of Animal Biology I Complutense University, Madrid, 1996	Duración River, Spain (released)	Fresh	402-bp
N1	<i>A. naccarii</i>	Aquatic Ecology Station, Seville	Sierra Nevada fish farm, Spain	Fresh	402-bp
N2	<i>A. naccarii</i>	Aquatic Ecology Station, Seville	Sierra Nevada fish farm, Spain	Fresh	402-bp
N3	<i>A. naccarii</i>	Aquatic Ecology Station, Seville	Sierra Nevada fish farm, Spain	Fresh	402-bp

samples were incubated again overnight at 55 °C. Standard phenol-dichloromethane 1:1 extractions were used to isolate DNA. DNA was precipitated in 1.6 ml of ice-cold 100 % EtOH and 0.1 vol. of 3 M sodium acetate (pH 6.8); incubated at -20 °C overnight; centrifuged at 16 000 g for 5 min at 4 °C; and washed in cold 70 % EtOH. Samples were air-dried and resuspended in 10-30 µl of ddH<sub>2</sub>O.

### PCR amplification

The PCR 402-bp fragments of the cytochrome *b* (cyt *b*) gene were amplified using universal primers (L14841; 5'-AAAAAGCTTCCATCCAACATCTCAGCATGATGAAA-3', H15149; 5'-AAACTGCAGCCCCTCAGAATGATATTTGTCCTCA-3', Kocher *et al.*, 1989) synthesized by Boehringer Mannheim (Spain). Tissues may degrade due to the storage process, making DNA extraction from archival specimens extremely difficult (France and Kocher, 1996; Rosenbaum *et al.*, 1997; Wirgin *et al.*, 1997). Therefore, we designed a specific internal primer (L14993; 5'-ATGACTAATCCGAAATATTC-3') to generate a 155 bp fragment in combination with H15149. This region was selected because of the high variability in it among *A. sturio*, *A. naccarii* and *A. baerii*.

Amplification was performed in 50 µl total volumes containing 1 unit of *Taq* I Polymerase (Biotools, Madrid, Spain), 5 µl of 10× reaction buffer (Biotools), 20 pmol of each primer, 2 mM MgCl<sub>2</sub>, 1 µl of dNTP mix (12.5 mM stock; Biotools), and ddH<sub>2</sub>O to volume. One microlitre of DNA was added to the PCR mix, and subjected to 35 cycles of amplification. DNA was amplified in a Perkin Elmer 9600 thermocycler using the following protocol: after a preliminary denaturation at 94 °C for 5 min, each cycle consisted of denaturation at 94 °C for 30 s, annealing at 44 °C for 30 s, and primer extension at 72 °C for 1 min. To improve the subsequent reamplification and sequencing, a final cycle was added, including an extension of 5 min at 72 °C. Negative controls (no DNA added) were run for each amplification. PCR products were electrophoretically separated in 1-1.5 % agarose gels with a molecular size ladder and stained in ethidium bromide solution to determine size and quality of fragments. Some PCR products from archived specimens were reamplified, in order to increase the quantity of DNA.

The amplification products were purified before sequencing, either with Qiaquick columns (Qiagen, Santa Clarita, CA, USA), or Biotools columns (Biotools, Madrid, Spain), following the manufacturer's instructions.

## Sequencing of PCR products

After purification, sequencing was performed in both directions by dideoxy sequencing (Sanger, Nicklen and Coulson, 1977) with the primers we used for PCR amplification. Sequencing was performed in an ABI PRISM 377 automated sequencer (Applied Biosystems, Foster City, USA), using four-dye fluorescent technology.

## The cyt *b* gene sequences

The following cyt *b* gene sequences from the related sturgeon species retrieved from GenBank were used for the phylogenetic study: *A. sturio* (accession no. AF006134), *A. baerii* (AF006123), *A. naccarii* (AF006150), *Acipenser sinensis* Gray, 1835 (AF006158), *Acipenser persicus* Borodin, 1897 (AF006156), *Acipenser oxyrinchus* Mitchill, 1815 (AF006154), *Acipenser gueldenstaedtii* Brandt & Ratzeberg, 1833 (AF006126), *Acipenser brevirostrum* LeSueur, 1818 (AF006124) and *Polyodon spathula* (Walbaum, 1792) (MTPSPCYTB).

## Sequence analysis

Sequences were analysed on an ABI PRISM 377 automated sequencer and data were edited with a multiple sequence editor program (ABI 1992). The sequences are deposited in GenBank under accession numbers AF217206-AF217209.

All DNA sequences were aligned using Clustal X (Thompson, Higgins and Gibson, 1994). Sequence data obtained for the long (402 bp) and short (155 bp) cyt *b* segments were analysed for distance and character-based variation using PHYLIP (version 3.5; Felsenstein, 1993). Phylogenetic trees were generated according to the maximum parsimony, MP (Felsenstein, 1983), maximum likelihood, ML (Felsenstein, 1981), and the distance-based neighbour-joining, NJ (Saitou and Nei, 1987) algorithms. For the transition/transversion bias, pairwise distances between nucleotide sequences were computed according to Kimura's two-parameters model (Kimura, 1980). The statistical significance of branching orders was assessed by the bootstrap resampling technique (100 replicates). Trees were rooted using as an outgroup *P. spathula*, a close relative of the Acipenseridae (Bemis, Findeis and Grande, 1997). A majority-rule consensus tree was constructed using the CONSENSE program from the PHYLIP package. The Kishino-Hasegawa-Templeton test (Templeton, 1983; Kishino and Hasegawa, 1989) was used to determine branch lengths from distance and ML test analyses.

## RESULTS

### Characteristics of the cyt *b* gene partial sequences

The long amplified fragment (402-bp) corresponds to the 5'-end of the cyt *b* gene. The short am-

ATGGCAAACA	TCCGAAAAAC	ACACCCACTA	CTTAAAATTA	TTAATGGAGC	50
ATTCATTGAC	CTCCCCACAC	CCTCCAACAT	CTCCGTGTGA	TGAAAYTTTG	100
GCTCACTCMT	AGGCCTCTGC	CTTGTSACAC	AAATCYTAAC	AGGACTATTT	150
CTTGCAATAC	AYTACACAGC	YGACATTTCA	ACAGCCTTCT	CCTCYRTYGC	200
CCACATCTGC	CGAGAYGTAA	ATTACGGATG	ACTAATCCGA	AATATTCATG	250
CAAACGGGGC	CTCYTTCTTC	TTYATYTGCT	TGTACCTTCA	CGTAGCACGA	300
GGYATR TACT	AYGGYTCATA	CCTCCAAAAA	GAAACCTGAA	AYATYGGRRGT	350
RATCCTYCTG	CTYCTCACCA	TAATAACCGC	CTTCGTGGGR	TATGTAYTRC	400
CC					402

Figure 1. Consensus sequence of a 402-bp cytochrome *b* fragment from *A. sturio*, *A. naccarii* and *A. baerii*. Numbers mark the variable positions among genotypes observed

Table II. Variable nucleotide positions of cytochrome *b* fragment sequences. Numbers refer to sequence position in figure 1

Genotypes	Sample	Variable nucleotides																									
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
STU	S1	?	?	?	?	?	?	?	?	?	?	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
STU	S2	C	C	G	T	T	C	C	G	C	T	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
STU	S3	C	C	G	T	T	C	C	G	C	T	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
STU	S4	C	C	G	T	T	C	C	G	C	T	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
STU	S5	C	C	G	T	T	C	C	G	C	T	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
STU	S6	?	?	?	?	?	?	?	?	?	?	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
STU	S7	?	?	?	?	?	?	?	?	?	?	T	C	T	C	A	C	C	C	C	G	G	T	C	G	T	A
BAE 1	B1	T	C	C	C	C	T	T	A	T	C	C	T	C	T	G	T	T	C	C	A	A	C	T	A	C	G
BAE 2	B2	T	A	C	C	C	T	T	A	T	C	C	T	C	T	G	T	T	C	C	A	A	C	T	A	C	G
BAE 1	B3	T	C	C	C	C	T	T	A	T	C	C	T	C	T	G	T	T	C	C	A	A	C	T	A	C	G
NAC	N1	T	C	C	C	C	T	T	G	T	C	C	T	C	T	A	T	T	T	T	A	A	C	T	A	C	G
NAC	N2	T	C	C	C	C	T	T	G	T	C	C	T	C	T	A	T	T	T	T	A	A	C	T	A	C	G
NAC	N3	T	C	C	C	C	T	T	G	T	C	C	T	C	T	A	T	T	T	T	A	A	C	T	A	C	G

plified fragment (155-bp) corresponds to positions from 247 to 402 of the *cyt b* gene of the white sturgeon, *Acipenser transmontanus* Richardson, 1836 (Brown *et al.*, 1989; GenBank accession no. X14944).

Twenty-six polymorphic sites were found on the 402-bp fragment among 10 specimens of *A. sturio*, *A. baerii*, and *A. naccarii* (figure 1). They showed a single base-pair change and the sequence variation was predominantly due to transitions ( $n = 24$ ). No deletions or insertions were observed. When we compared the 155-bp fragment obtained from S1, S6, and S7 with the other samples, we found 16 changes due to transitions in the sequences (table II). Both regions studied had a low G content (average 16.2 %) and almost equal A, C, and T contents (average 26.7, 29.3, and 27.8 %, respectively). The base composition of this *cyt b* region in *Acipenser* was similar to that found in teleosts (rep-

resentatives of the order Perciformes) (Cantatore *et al.*, 1994).

### Sequence variation and diversity of the *cyt b* genotypes

Four *cyt b* genotypes were identified in *A. sturio* (STU), *A. naccarii* (NAC), and *A. baerii* (BAE-1 and BAE-2) (table II). The sequence from the *A. sturio* specimen EBD 8174 was identical to those from sturgeon representing the Gironde River, Adriatic, and North Sea populations. However, at position 126 we found G in all *A. sturio* sequences, whereas Birstein and DeSalle (1998) identified T in the sequence from a North Sea individual. We did not observe any variable site within this fragment in *A. naccarii*, and we found a substitution (position 109) in one *A. baerii*.

Table III. Genetic distances among 245-bp sequences analysed following Kimura's two-parameter model. Cells in bold characters show the genetic distance analysed in the 155-bp sequence

	STU	BAE-1	BAE-2	NAC	<i>Acipenser sinensis</i>	<i>Acipenser persicus</i>	<i>Acipenser naccarii</i>	<i>Acipenser gueldenstaedtii</i>	<i>Acipenser brevirostrum</i>	<i>Acipenser oxyrinchus</i>	<i>Acipenser sturio</i>	<i>Polyodon spathula</i>
STU	0.00											
BAE-1	6.91	0.00										
BAE-2	7.43	0.41	0.00									
NAC	6.91	1.65	2.09	0.00								
<i>A. sinensis</i>	10.80	9.78	10.35	10.75	0.00							
<i>A. persicus</i>	6.91	1.65	2.09	1.65	10.75	0.00						
<i>A. naccarii</i>	7.83	2.50	2.95	0.82	11.73	2.50	0.00					
<i>A. gueldenstaedtii</i>	6.91	0.00	0.41	1.65	9.78	1.65	2.50	0.00				
<i>A. brevirostrum</i>	6.91	0.82	1.25	2.50	10.75	2.50	3.35	0.82	0.00			
<i>A. oxyrinchus</i>	4.23	9.67	10.23	8.73	12.73	8.73	8.73	9.67	10.62	0.00		
<i>A. sturio</i>	0.41	6.89	7.40	6.89	10.75	6.89	7.80	6.89	6.89	4.22	0.00	
<i>P. spathula</i>	12.95	12.89	13.52	12.89	8.36	12.89	13.92	12.89	13.92	14.96	12.89	0.00

Pairwise estimates of sequence divergence in the 402-bp fragment of the *cyt b* gene among haplotypes of *A. sturio*, *A. baerii* and *A. naccarii* were 0.41–7.43 % (table III). Evidently, the American Atlantic sturgeon *A. oxyrinchus* is the closest relative of *A. sturio*: we observed a 4.2 % divergence between the *cyt b* sequences of these species. *P. spathula*, which we used as an outgroup, presented the highest sequence divergence with *A. sturio* (12.9 %). *A. baerii* and *A. naccarii* showed a similar divergence from *A. sturio* (6.9 and 7.4 %, respectively).

The 402-bp fragment of the *cyt b* gene sequence was aligned with those of the related sturgeon species retrieved from GenBank. A phylogenetic analysis of this sequence data was performed to determine the position of the samples. The nucleotide sequence alignment consensus for all taxa was 245 bp long. Figure 2 shows the consensus tree built using the NJ method with a 7.17 ts:tv weighting assumption; *P. spathula* was used as an outgroup. The MP and ML analyses resulted in a similar and congruent tree after collapsing poorly




Figure 2. Estimated phylogenetic tree of 245-bp cytochrome *b* sequences. The percentages of bootstrap replicates supporting the clades are indicated at the branch points, based on (from left to right) neighbour-joining, maximum likelihood and equally-weighted parsimony methods. Bootstrap values < 50 % are not shown. Branch lengths are proportional to the estimated mean number of substitutions per site. GenBank species names are underlined






Figure 3. Estimated phylogenetic tree of 105-bp cytochrome *b* sequences according to the neighbour-joining method Kimura's distance. Bootstrap supports higher than 50 % are shown. Branch lengths are proportional to the estimated mean number of substitutions per site. GenBank species names are underlined

supported bootstrap values. The position of *A. sinensis* was not clearly resolved. Two clusters, A and B, can be clearly identified. Cluster A contained *A. persicus* (I), a clade of the *A. naccarii* specimens (II), and a clade which included *A. brevirostrum*, *A. gueldenstaedtii* and the *A. baerii* specimens (III). *A. persicus* was a sister species to (II) and (III). The clade (II) included all *A. naccarii* specimens, archival and alive. Its monophyly was well-supported (88 % for the NJ tree, 60 % ML tree, and 90 % for the MP tree). The *A. baerii* specimens analysed in the present paper, and the *A. baerii*, *A. brevirostrum*, and *A. gueldenstaedtii* specimens studied by Birstein

and DeSalle (1998) (data from GenBank) also formed a well-supported clade. The second cluster, B, contained *A. oxyrinchus* and *A. sturio*.

The second phylogenetic tree was built using the 155-bp cyt *b* gene fragments obtained from *A. sturio* museum specimens originating from the Iberian Peninsula, Gironde River, North Sea, and the Adriatic (figure 3). The same overall pattern was observed for this short sequence as with the previous 245-bp segment. Sequences from all *A. sturio* samples, including that retrieved from GenBank, were clustered together, forming a monophyletic branch.

## DISCUSSION

### DNA from archival specimens

Currently, museum specimens are the only available material for *A. sturio* from many populations or from a geographical range of a population that has almost disappeared. Recent advances in molecular techniques offer a potential for DNA extraction from a diverse range of materials previously considered intractable (France and Kocher, 1996; Habelberg, Sykes and Hedges, 1989; Rosenbaum *et al.*, 1997; Pääbo, 1989; Wirgin *et al.*, 1997; see DeSalle and Bonwich, 1996, for a review). In our study the best yield in PCR products from archival samples was when we followed the protocol of France and Kocher (1996). However, only a short 155-bp fragment of the *cyt b* gene was successfully amplified. The tree based on the 105-bp sequences showed that a short sequence is suitable for discerning among the species studied: the nodes had good support bootstrap values (figure 3). We agree with Ludwig *et al.* (2000) that since *A. sturio* has almost disappeared in the wild, molecular analyses of specimens representing different populations in museum collections are crucial for any recovery plan for this species.

### Phylogenetic implications

Clusters A and B in our trees (figures 2 and 3) generally support the phylogenetic relationships presented in Birstein and DeSalle (1998) for a longer region (650-bp) of the *cyt b* gene and for combined data for this region, plus a 350-bp fragment of the 16S rDNA and a 150-bp fragment of the 12S rDNA (figures 1 and 3 in that paper). The different position of *A. sinensis* in our tree, compared to the trees in Birstein and DeSalle (1998), was evidently caused by the high divergence of this *cyt b* gene fragment in *A. sinensis* from the other *Acipenser* species included in our study.

The relationships of species within clades of cluster A are basically congruent with the earlier phylogenetic studies. The position of the North American shortnose sturgeon of the Atlantic coast, *A. brevirostrum*, in our tree point to a close relatedness to the Russian sturgeon *A. gueldenstaedtii* inhabiting the Ponto-Caspian basin (Vlasenko *et al.*, 1989), and with the Siberian sturgeon *A. baerii*, liv-

ing in Siberian rivers and Lake Baikal (Ruban, 1997). Morphologically *A. gueldenstaedtii* is similar to *A. persicus* (Vlasenko *et al.*, 1989) and *A. naccarii* (Tortonese, 1989), and it is considered to be related to *A. baerii* (Sokolov and Vasil'ev, 1989). Our results showed that *A. gueldenstaedtii* appears to be more closely related to *A. baerii* and *A. brevirostrum* than to *A. naccarii* and *A. persicus*. *A. persicus* is a sister species to the *A. baerii*, *A. gueldenstaedtii* and *A. brevirostrum* cluster, and to the *A. naccarii* clade. Conversely, Tagliavini *et al.* (1999) postulated a close relationship between *A. naccarii* and *A. gueldenstaedtii* and other clusters which includes *A. brevirostrum* and *A. baerii*.

Although Birstein and DeSalle (1998) used a much longer region of the *cyt b* gene (650 bp), their data showed that *A. gueldenstaedtii*, *A. persicus*, *A. baerii*, *A. naccarii*, and *A. brevirostrum* form an unresolved clade (figure 1 in their paper). They succeeded in placing *A. brevirostrum* outside this clade only when they used combined data for three genes, which contained a 1150-bp-long sequence. In this analysis, *A. brevirostrum* was basal to the Ponto-Caspian and some Asian sturgeons (figure 3 in Birstein and DeSalle, 1998). A more detailed study of the relationships among Eurasian species of the same clade included a longer region of the *cyt b* gene (850-bp), a partial sequence of the ND5 gene (643-bp), and a fragment of the control region of mitochondrial DNA (725-bp), or approximately 2.3 kb bp (Birstein, Doukakis and DeSalle, in press). The traditionally recognised *A. gueldenstaedtii* in fact appeared to consist of two genetic forms, a "typical" and a cryptic one. The "typical" form is phylogenetically affiliated with *A. persicus* and *A. naccarii*. These data show that *A. persicus* is not a separate species; rather, it is conspecific with *A. gueldenstaedtii*. *A. naccarii* was diagnosed by only one G in position 164 of the control region. The cryptic form of *A. gueldenstaedtii* is very close genetically to *A. baerii*: it differed from *A. baerii* only by C-T transition at position 432 of the control sequence.

This example of complex species relationships found among a group of Eurasian sturgeons illustrates that reliable phylogenetic data can be obtained only if long DNA sequences are included in a study. Such results as those published by Tagliavini *et al.* (1999) can be considered only preliminary: these authors used very short sequences in their research and studied a small number of



sturgeon species. The latter leads to an inadequate picture of species relationships due to the problem of underdiagnosis of phylogenetic lineages (Davis and Nixon, 1992). In general, the low genetic variability shown by acipenseriforms (Birstein, Hanner and DeSalle, 1997) makes a study of phylogenetic relationships within this order complicated.

#### ***A. sturio*/*A. oxyrinchus* cluster and *A. sturio* intraspecies variability**

The grouping of *A. sturio* with *A. oxyrinchus* (figures 2 and 3) is congruent with previous molecular phylogenetic results, and supports the hypothesis that these species might be direct descendants of ancestral forms of *Acipenser* (Birstein and DeSalle, 1998). The present molecular study has not detected genetic difference among specimens of *A. sturio* from the Iberian Peninsula, Gironde River, North Sea, and Adriatic. Likewise, Ludwig and Kirschbaum (1998) did not find variable sites in a 400-bp fragment of the 12S ribosomal DNA when they compared *A. sturio* specimens from the North Sea and Gironde River. However, Birstein and DeSalle (1998) obtained six nucleotide changes in a 600-bp fragment of cyt *b* in two specimens of *A. sturio*, suggesting intra-species forms of *A. sturio* (see Birstein and Doukakis, 2000, for further discussion).

#### **The specimen EBD 8174**

Garrido-Ramos *et al.* (1997) found a HindIII satellite (st) DNA in the genomes of two specimens of *A. naccarii* from a fish farm that was absent in two specimens of *A. sturio* from the Gironde River. The HindIII stDNA family was then considered a species-specific nuclear DNA marker (see Ruiz-Rejón *et al.*, 2000). Garrido-Ramos *et al.* (1997) attributed specimen EBD 8174 to *A. naccarii*, based on the presence of the HindIII sequences in the DNA extracted from this specimen. The present study enabled us to positively identify specimen EBD 8174 as *A. sturio*. Our conclusion supports the results of previous morphometric and meristic studies of this specimen (Elvira and Almodóvar, 1999, 2000; Rincón, 2000).

## **CONCLUSIONS**

Our preliminary results did not show genetic differences between *A. sturio* individuals from the Iberian Peninsula, Gironde River, Adriatic, and North Sea. We identified specimen EBD 8174 from the Guadalquivir River as *A. sturio*. The comparison of the cyt *b* gene sequences among *A. sturio*, *A. naccarii*, and *A. baerii* showed 22-24 fixed nucleotide changes among species in that region. The analysis of our data set, together with that retrieved from GenBank, supports the hypothesis of Birstein and DeSalle (1998) that *A. oxyrinchus* is the only sturgeon species closely related to *A. sturio*. Since *A. sturio* has almost disappeared in the wild, the molecular analyses of specimens representing different populations of this species in museum collections are specially important in terms of recovery plans for this species.

## **ACKNOWLEDGEMENTS**

We are grateful to B. Elvira for providing us with samples used in the study and for helpful discussions. We thank M. Lepage and E. Rochard for sturgeon samples from the Gironde River. We also thank P. Rubio and the Department of Cellular and Molecular Biology (University of Alcalá, Madrid) for technical assistance in sequencing. We are also grateful to G. G. Nicola for reviewing the manuscript. This work received support from the Agricultural Research Institute of Madrid (IMIA).

## **REFERENCES**

- Almaça, C. 1988. On the sturgeon, *Acipenser sturio*, in the Portuguese rivers and sea. *Folia Zoologica* 37: 183-191.
- Bemis, W. E., E. K. Findeis and L. Grande. 1997. An overview of Acipenseriformes. In: *Sturgeon Biodiversity and Conservation*. V. J. Birstein, J. R. Waldman and W. E. Bemis (eds.): 25-71. Kluwer Academic Publishing. Dordrecht, Netherlands.
- Birstein, V. J. and W. E. Bemis. 1997. How many species are there within the genus *Acipenser*? In: *Sturgeon Biodiversity and Conservation*. V. J. Birstein, J. R. Waldman and W. E. Bemis (eds.): 157-163. Kluwer Academic Publishing. Dordrecht, Netherlands.
- Birstein, V. J., J. Betts and R. DeSalle. 1998. Molecular identification of *Acipenser sturio* specimens: a warning note for recovery plans. *Biological Conservation* 84: 97-101.

- Birstein, V. J. and R. DeSalle. 1998. Molecular phylogeny of Acipenserinae. *Molecular Phylogenetics and Evolution* 9 (1): 141-155.
- Birstein, V. J. and P. Doukakis. 2000. Molecular analysis of *Acipenser sturio* L., 1758 and *Acipenser oxyrinchus* Mitchell, 1815: A review. In: *Symposium on Conservation of the Atlantic Sturgeon Acipenser sturio L., 1758 in Europe*. Boletín. Instituto Español de Oceanografía 16 (1-4): 61-73.
- Birstein, V. J., P. Doukakis and R. DeSalle. (In press). Caviar species identification and the polyphyletic nature of Russian sturgeon: forensic and evolutionary implications. *Conservation Genetics*.
- Birstein, V. J., R. Hanner and R. DeSalle. 1997. Phylogeny of the Acipenseriformes: cytogenetic and molecular approaches. In: *Sturgeon Biodiversity and Conservation*. V. J. Birstein, J. R. Waldman and W. E. Bemis (eds.): 127-155. Kluwer Academic Publishing. Dordrecht, Netherlands.
- Bonaparte, C. 1836. *Iconografia della fauna italiana per le quattro classi degli animali vertebrati*. Tomo III. Pesci. Roma. Fasc. 15-18: 80-93.
- Brown, J. R., T. L. Gilbert, D. J. Woebel, P. J. O'Hara, N. E. Buroker, A. T. Beckenbach and M. J. Smith. 1989. Nucleotide sequence of the apocytochrome B gene in white sturgeon mitochondrial DNA. *Nucleic Acids Research* 17: p. 4389.
- Cantatore, P., M. Roberti, G. Pesole, A. Ludovico, F. Milella, M. Gadaleta and C. Saccone. 1994. Evolutionary analysis of cytochrome b sequences in some Perciformes: evidence for a slower rate of evolution than in mammals. *Journal of Molecular Evolution* 39: 589-597.
- Davis, J. I. and K. C. Nixon. 1992. Populations, genetic variation, and the determination of phylogenetic species. *Systematic Biology* 41: 421-435.
- DeSalle, R. and E. Bonwich. 1996. DNA isolation, manipulation and characterization from old tissues. In: *Genetic Engineering*. J. Setlow (ed.) 18: 13-22. Plenum. New York.
- Doadrio, I., B. Elvira and Y. Bernat. 1991. *Peces continentales españoles. Inventario y clasificación de zonas fluviales*. Icona, Colección Técnica. Madrid: 221 pp.
- Doukakis, P., V. J. Birstein, R. DeSalle, A. N. Ludwig, A. Ludwig, A. Machordom, A. Almodóvar and B. Elvira. 2000. Failure to confirm previous identification of two putative museum specimens of the Atlantic sturgeon, *Acipenser sturio*, as the Adriatic sturgeon, *A. naccarii*. *Marine Biology* 136: 373-377.
- Elvira, B. and A. Almodóvar. 1993. Notice about the survival of sturgeon (*Acipenser sturio* L., 1758) in the Guadalquivir estuary (S.W. Spain). *Archiv für Hydrobiologie* 129 (2): 253-255.
- Elvira, B. and A. Almodóvar. 1999. A morphological study of native sturgeon *Acipenser sturio* in Spain, and recent records of exotic Siberian sturgeon *A. baerii*. *Journal of Applied Ichthyology* 15 (4-5): 278-279.
- Elvira, B. and A. Almodóvar. 2000. Morphology and taxonomy of the Atlantic sturgeon *Acipenser sturio* from Spain. *Folia Zoologica* 49 (3): 221-230.
- Elvira, B., A. Almodóvar and J. Lobón-Cerviá. 1991a. Recorded distribution of sturgeon (*Acipenser sturio* L., 1758) in the Iberian Peninsula and actual status in Spanish waters. *Archiv für Hydrobiologie* 121 (2): 253-258.
- Elvira, B., A. Almodóvar and J. Lobón-Cerviá. 1991b. Sturgeon (*Acipenser sturio* L., 1758) in Spain. The population of the river Guadalquivir: a case history and a claim for a restoration programme. In: *Acipenser*. P. Williot (ed.): 337-347. Cemagref. Bordeaux.
- Felsenstein, J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution* 36: 182-198.
- Felsenstein, J. 1983. Parsimony in systematics: biological and statistical issues. *Annual Review of Ecology and Systematics* 14: 313-333.
- Felsenstein, J. 1993. *PHYLIP: Phylogeny Inference Package. Version 3.5*. University of Washington. Seattle, WA.
- France, S. C. and T. D. Kocher. 1996. DNA sequencing of formalin-fixed crustaceans from archival research collections. *Marine Biology and Biotechnology* 5 (4): 304-313.
- Garrido-Ramos, M. A., M. C. Soriguer, R. de la Herrán, M. Jamilena, C. Ruiz-Rejón, A. Domezain, J. A. Hernando and M. Ruiz-Rejón. 1997. Morphometric and genetic analysis as proof of the existence of two sturgeon species in the Guadalquivir River. *Marine Biology* 129: 33-39.
- Habelberg, E., B. Sykes and R. Hedges. 1989. Ancient bone DNA amplified. *Nature* 342: 485.
- Holčík, J., R. Kinzelbach, L. I. Sokolov and V. P. Vasil'ev. 1989. *Acipenser sturio* Linnaeus, 1758. In: *The Freshwater Fishes of Europe. General Introduction to Fishes, Acipenseriformes*. J. Holčík (ed.) 1 II: 367-394. AULA-Verlag, Wiesbaden, Germany.
- Kimura, M. 1980. A simple model for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16: 11-120.
- Kishino, H. and M. Hasegawa. 1989. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order of the Hominoidea. *Journal of Molecular Evolution* 29: 170-179.
- Kocher, T. D., W. K. Thomas, A. Meyer, S. V. Edwards, S. Pääbo, F. X. Villablanca and A. C. Wilson. 1989. Dynamics of mitochondrial DNA evolution in animals: Amplification and sequencing with conserved primers. *Proceedings of National Academy of Sciences (USA)* 86: 6196-6200.
- Ludwig, A. N., I. Jenneckens, L. Debus, A. Ludwig, J. Becker, and F. Kirschbaum. 2000. Genetic analyses of archival specimens of the Atlantic sturgeon *Acipenser sturio* L., 1758. In: *Symposium on Conservation of the Atlantic Sturgeon Acipenser sturio L., 1758 in Europe*. Boletín. Instituto Español de Oceanografía 16 (1-4): 181-190.
- Ludwig, B. and F. Kirschbaum. 1998. Comparison of mitochondrial sequences of the Adriatic and the European sturgeon. *Journal of Fish Biology* 52: 1289-1291.
- Pääbo, S. 1989. Ancient DNA: extraction, characterization, molecular cloning and enzymatic amplification. *Proceedings of National Academy of Sciences (USA)* 86: 1939-1943.
- Pereira, N. C. 1995. The freshwater fishes of the Iberian Peninsula. *Archivos Museu Bocage. Nova Serie* 2: 473-538.

- Rincón, P. A. 2000. Big fish, small fish: still the same species. Lack of morphometric evidence of the existence of two sturgeon species in the Guadalquivir River. *Marine Biology* 136 (4): 715-723.
- Rosenbaum, H. C., M. G. Egan, P. J. Clapham, R. L. Brownell and R. DeSalle. 1997. An effective method for isolating DNA from historical specimens of baleen. *Molecular Ecology* 6: 677-681.
- Ruban, G. I. 1997. Species structure, contemporary distribution and status of the Siberian sturgeon, *Acipenser baerii*. In: *Sturgeon Biodiversity and Conservation*. V. J. Birstein, J. R. Waldman and W. E. Bemis (eds.): 221-230. Kluwer Academic Publishing, Dordrecht, Netherlands.
- Ruiz-Rejón, M., R. de la Herrán, C. Ruiz-Rejón and M. A. Garrido-Ramos. 2000. Genetic characterization of *Acipenser sturio* L., 1758 in relation to other sturgeon species using satellite DNA. In: *Symposium on Conservation of the Atlantic Sturgeon Acipenser sturio* L., 1758 in Europe. *Boletín. Instituto Español de Oceanografía* 16 (1-4): 231-236.
- Saitou, N. and M. Nei. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4: 406-425.
- Sanger, F., S. Nicklen and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proceedings of National Academy of Sciences (USA)* 74: 5463-5467.
- Shiozawa, D. K., J. Kudo, R. P. Evans, S. R. Woodward and R. N. Williams. 1992. DNA extraction for preserved trout tissues. *Great Basin Naturalist* 52: 29-34.
- Sokolov, L. I. and V. P. Vasil'ev. 1989. *Acipenser baeri* Brandt, 1869. In: *The Freshwater Fishes of Europe. General Introduction to Fishes, Acipenseriformes*. J. Holčík (ed.) 1 II: 263-284. AULA-Verlag, Wiesbaden, Germany.
- Tagliavini, J., F. Conterio, G. Gandolfi and F. Fontana. 1999. Mitochondrial DNA sequences of six sturgeon species and phylogenetic relationships within Acipenseridae. *Journal of Applied Ichthyology* 15 (4-5): 17-22.
- Templeton, A. R. 1983. Phylogenetic inference from restriction endonuclease cleavage site maps with particular reference to the humans and apes. *Evolution* 37: 221-244.
- Thompson, J. D., D. G. Higgins and T. J. Gibson. 1994. CLUSTAL X: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673-4680.
- Tortonese, E. 1989. *Acipenser naccarii* Bonaparte, 1836. In: *The Freshwater Fishes of Europe. General Introduction to Fishes, Acipenseriformes*. J. Holčík (ed.) 1 II: 285-293. AULA-Verlag, Wiesbaden, Germany.
- Vlasenko, A. D., A. V. Pavlov, L. I. Sokolov and V. P. Vasil'ev. 1989. *Acipenser gueldenstaedti* Brandt, 1833. In: *The Freshwater Fishes of Europe. General Introduction to Fishes, Acipenseriformes*. J. Holčík (ed.) 1 II: 294-344. AULA-Verlag, Wiesbaden, Germany.
- Wirgin, I., L. Maceda, J. Stabile and C. Mesing. 1997. An evaluation of introgression of Atlantic coast striped bass mitochondrial DNA in a Gulf of Mexico population using formalin-preserved museum collections. *Molecular Ecology* 6: 907-916.
- Wirgin, I. I., J. E. Stabile and J. R. Waldman. 1997. Molecular analysis in the conservation of sturgeons and paddlefish. In: *Sturgeon Biodiversity and Conservation*. V. J. Birstein, J. R. Waldman and W. E. Bemis (eds.): 385-398. Kluwer Academic Publishing, Dordrecht, Netherlands.