





## pBR322 DNA/BsuRI (HaeIII) Marker (Ref. 31.003)

Concentration: 0.5 mg/ml (50 µg)

Store at -20°C

### Description

The pBR322 DNA is completely digested with BsuRI to yield bands ranging from 8 bp to 587 bp, suitable for use as molecular weight standards for agarose and polyacrylamide gels. The marker is composed to 22 purified individual DNA fragments (in base pairs): 587, 540, 502, 458, 434, 267, 234, 213, 192, 184, 124, 123, 104, 89, 80, 64, 57, 51, 21\*, 18\*, 11\* and 8\*.



\*The shortest fragments (21, 18, 11 and 8 bp) are not visible in standard electrophoresis

\*\*The 587, 540 and 458 bp fragments migrate anomalously on polyacrylamide gels

### Storage buffer (TE buffer)

10 mM Tris-HCl (pH 7.6), 1 mM EDTA

### Storage temperature

Store at -20°C. For frequent use divide in aliquots to avoid multiple freeze/thaw cycles, or store at 4°C in the presence of loading buffer (stable for 4 months).

## Protocol

1- Prepare loading mixture (for the 5 mm gel lane\*):

	Agarose Gels	Polyacrylamide Gels
• DNA Marker (0.5-1 µg)	1 µl	2 µl
• 6X Loading Dye	1 µl	0.5 µl
• Distilled water	4 µl	0.5 µl
Final volume	6 µl	3 µl

2- Mix gently

**3- Do not heat**

4- Load onto the gel

5- Visualise DNA by staining with ethidium bromide or with SYBR® Green I

\*The mixture should be scaled up or down, depending on the width of the gel. Use 0.1-0.2 µg of DNA marker/mm of lane.

The pBR322 DNA/BsuRI Marker was not designed for precise quantification of DNA mass, but can be used for semi-quantification (see Table 1). For quantification, adjust the concentration of the sample to equalize it approximately with the amount of DNA in the nearest band of the ladder.

Table 1. Percentage and mass of individual fragments (for 0.5 µg pBR322 DNA/BsuRI Marker)

Fragment	Size	%	mass (ng/0.5µg)
1	587	13.5	67.3
2	540	12.4	61.9
3	502	11.5	57.6
4	458	10.5	52.5
5	434	10.0	49.8
6	267	6.1	30.6
7	234	5.4	26.8
8	213	4.9	24.4
9	192	4.4	22.0
10	184	4.2	21.1
11	124	2.8	14.2
12	123	2.8	14.1
13	104	2.4	11.9
14	89	2.0	10.2
15	80	1.8	9.2

### Notice to users

Research and *in vitro* use only.